

Molecular Otolaryngology & Renal Research Laboratories For test inquiries please call: 319-335-6623 • For billing inquiries call: 319-467-1647

Fax: 319-353-5869 or 319-335-9890

Email: morl@uiowa.edu • https://morl.lab.uiowa.edu

CLIA: 16D0966193

MORL - Kidney Testing Requisition Form

REFERRING LABORATORY USE ONLY : please complete below section						F	FOR MORL USE ONLY	<i>(</i> :			
Requisition Date:	Comp	leted by:		Accn#:			٢	MORL Case #:			
Collection: Blood Dat	te/Time:	#Tubes	: Serum	Date/Time:	:	#Tubes:	Plasr	ma Date/Time:		#Tubes:	
Part A) Patient	Information or	ID Sticke	er <u>(Required)</u>		Part A	1) Patient Demo	grapł	hic Information	(Required	<u>1)</u>	
Name:					Ethnicit	t y : 🗌 Hispanio	с	🗌 Not Hisp	anic		
Last		First			Rac	e : 🗌 White		Black or	African An	nerican	
DOB: /	day /year	Sex: (please of	□ Male □ Fe check sex assigned	male d at birth)		Asian		America	n Indian/A	laska Na	ative
Height:						Native H	lawaiia	an or Other Pacif	ic Islander		
						🗌 More Th	ian On	ne Race			
MRN:				Part C) Payment Information (<i>Required</i>)							
Part B) Reporti	ng Information	(<u>Required</u>	<u>d)</u>		Institutional billing or payment by Visa or MasterCard is accepted.						
Health Care Provi	ider:				***The	MORL will <u>NOT</u> s	submi	t to insurance.			
E-mail Address:					Billing Contact:						
Institution:					Institution:						
Street Address:				Street Address:							
City:	State:		Zip:		City:		State	2:	Zip:		
Phone: ()		FAX: ()		Phone:	()		FAX: ()			
lf y	ou or your pation	ent would	like to pay by	credit ca	ard, pleas	se contact us at <u>n</u>	norl@	uiowa.edu for in	structions		
Part D) Pertinent Clinical Information (<u>Required</u>) – Complete the section below											
Diagnosis:	aHUS: Trigger	? 🗌 No [Yes (if <u>yes</u> , de	escribe trigg	ger, eg. BM	T, pregnancy, pneumo	ococcal)):			
						Other (comp					
Family history of renal disease? No Yes (if <u>yes</u> , please provide details in comment & attach a pedigree if available)											
Disease History Date of symptom onset:					Specimen Inform	ation:	<u>.</u>				
Renal Biopsy:						Was specimen dra	wn	Duran I. I.		_	
Hematuria:	Yes 🗌 No	•				pre or post:		Procedure dat	e: Pre	Post	N/A
Diarrhea:	Yes No					Eculizumab:					
Schistocytes:	Yes 🗌 No					PLEX (*affects serol	logies):				
Current Lab Value				—		Renal Tx:			 		
Hg/Hct:	<u>Value</u>	<u>No</u>	rmal Range	<u>Test [</u>	<u>Date</u>				LI		
Haptoglobin:						BMT (*affects gene	tics):				
Platelets:						Liver Tx:					
sCr/BUN:											
eGFR:						Comments:					
LDH:											
- uProt/uCr:											
Urine Blood:											
C3 Level:											
C4 Level:											
ADAMTS13:											



Molecular Otolaryngology & Renal Research Laboratories For test inquiries please call: 319-335-6623 • For billing inquiries call: 319-467-1647

Fax: 319-353-5869 or 319-335-9890

Email: morl@uiowa.edu • https://morl.lab.uiowa.edu CLIA: 16D0966193

DOB: MRN: Patient Name: Please see page 3 for sample handling requirements - No Weekend Deliveries **Functional Testing Panels Requested Genetic Testing Requested** C3 Glomerulopathy Complement Panel (C3G-CP) Genetic Renal Panel (DNA test for TTP, aHUS, HUS, DDD, C3G and (Serologies for complement-mediated renal diseases) other complement diseases) (CH50, APFA, C3b Deposition Assay, FHAA, FBAA, Fluid (CFH, CFI, MCP, CFB, CFHR5, C3, THBD, ADAMTS13, PLG, Phase Activity Assay-IFE, Nephritic Factors (C3Nef-C3CSA, DGKE, G6PD, MMACHC, WT1 and MLPA) C5Nef-C3CSAP, C4Nef), C3, C3c, C4, FB, Ba, Bb, FD, • 3-5 cc EDTA whole blood (room temp or refrigerated) Properdin, C5, Soluble C5b-9, FH and FI levels) Saliva: DNA Genotek, ORAgene (1 saliva vial) OR ORAcollect • 2 mL frozen serum (at least 4 sponges) 2 mL frozen EDTA plasma OR 5 μg DNA, minimum concentration 50 ng/μl aHUS (complement-mediated TMA) Panel (aHUS-FP) \square MLPA Testing ONLY (screening for copy number variations in the CFH-(CH50, APFA, C3b Deposition, FHAA, FBAA, Fluid Phase CFHR5 genomic region) Activity-IFE, C3, C3c, C4, FB, Ba, Bb, FD, Properdin, C5, • 3-5 cc EDTA whole blood (room temp or refrigerated) Soluble C5b-9, FH and FI levels) • Saliva: DNA Genotek, ORAgene (1 saliva vial) OR ORAcollect • 2 mL frozen serum (at least 4 sponges) • 2 mL frozen EDTA plasma OR 5 μg DNA, minimum concentration 50 ng/μl \square Autoantibody Panel (AAP) (FHAA, FBAA, Fluid Phase Activity-IFE, Nephritic Factors \square Familial Testing (site specific analysis to screen for variants previously (C3Nef-C3CSA, C5Nef-C3CSAP, C4Nef) identified in a family member) • 2 mL frozen serum Familial Testing Details: Gene/s: **Complement Biomarker Panel (CBP)** (C3*, C3c, C4*, FB, Ba, Bb, C5, FD, Properdin, Soluble C5b-9, MORL ID# or Variant/s: _____ FH and FI levels) • 2 mL frozen serum* **Relationship to** previously tested person: _____ 2 mL frozen EDTA plasma \square **Complement Pathway Activity Panel (CPAP)** If you are interested in ordering Custom Testing please contact (CH50, APFA, C3b Deposition Assay) Amy Weaver at 319-335-6623 or amy-weaver@uiowa.edu • 2 mL frozen serum Important Information for ALL Requests a La Carte Testing Requested Autoantibody Tests – 1 mL Frozen Serum All serum and plasma samples MUST be processed and frozen down FH autoantibody (FHAA) FB autoantibody (FBAA) \square to -80° C immediately after collection (please see instructions on page 3). Sample type must be clearly labeled (either serum or Fluid Phase Activity (IFE) C3Nef (C3CSA) \square plasma) and shipped out overnight on at least 5 lb dry ice **C5Nef (C3CSAP)** C4Nef (Monday – Thursday). Biomarker Tests – 1 mL Frozen EDTA Plasma If samples arrive thawed they will be **REJECTED**. **C3 Level** (1 mL frozen serum) C3c Level \square **No Weekend Deliveries C4 Level** (1 mL frozen serum) □ FD Level \square FB Level Ba Level Bb Level \square **Molecular Otolaryngology & Renal Research Laboratories Properdin Level** □ Soluble C5b-9 \square C5 Level For test inquiries please call: 319-335-6623 • Fax: 319-353-5869 For billing inquiries please call: 319-335-6653 • Fax: 319-353-5869 FI Level FH Level Email: morl@uiowa.edu • https://morl.lab.uiowa.edu **Complement Pathway Function Tests – 1 mL Frozen Serum C3b** Deposition Assay Ship to: CH50 \square **Dr. Richard Smith** \square **APFA** (Alternative Pathway Functional Assay) Molecular Otolaryngology & Renal Research Laboratories The University of Iowa 285 Newton Rd., 5270 CBRB ADAMTS-13 Tests – 1 mL Frozen Citrate Plasma Iowa City, IA 52242-1078 □ ADAMTS-13 Activity Monday — Friday ONLY – No Weekend Deliveries ADAMTS-13 Activity with reflex to Inhibitor (when activity <25%) \square



Molecular Otolaryngology & Renal Research Laboratories For test inquiries please call: 319-335-6623 • For billing inquiries call: 319-467-1647

Fax: 319-353-5869 or 319-335-9890

Email: morl@uiowa.edu • https://morl.lab.uiowa.edu

CLIA: 16D0966193

Molecular Otolaryngology & Renal Research Laboratories Sample Requirements

Questions? Contact Amy Weaver at 319-335-6623 or amy-weaver@uiowa.edu Genetic Renal Panel/MLPA/DNA Testing Sample Requirements: 3-5 cc. EDTA whole blood – MORL not responsible for broken tubes Saliva: DNA Genotek, ORAgene (1 saliva vial) OR ORAcollect (at least 4 sponges) 5 µg DNA, minimum concentration 50ng/µl (A260/A280 1.8-2) resuspended in 0.1mM EDTA (10mM Tris HCl, 0.1mM EDTA, pH 8, Teknova Cat# T0220) *Please note: blood samples drawn from a bone marrow transplant patient will result in genetic results for the donor rather than the patient, buccal swab samples are recommended. Overnight delivery, Room temperature or refrigerated (DO NOT FREEZE WHOLE BLOOD) Samples are accepted Monday-Friday. Samples may be refrigerated if delivery is delayed (stability – 1 week) Serum (FH & FB autoantibody, APFA, C3b Deposition Assay, CH50, Fluid Phase Activity, C3Nef, C5Nef, C4Nef, C3 and C4 levels) Collection Protocol (minimum volume 2ml) *PLEX treatments will affect serum tests, please wait ~14 days after PLEX to draw samples: 1. Follow standard phlebotomy techniques to collect at least 6 cc of whole blood drawn in a red-top vacutainer tube. Note: Serum separators with "clot activators" should not be used for the serum samples. 2. Allow the blood in the red-top tube to clot at room temperature for 30 minutes. 3. Centrifuge the clotted blood at room temperature (1000 x g for 10 minutes). 4. Label "Serum" or "Red-top" on clean screw top-tube (s). Pipette cell-free supernatant (at least 2 mL) to each labeled tube (s). 5. 6. Place the tube immediately at -80°C (or on dry ice). Sample must remain deep frozen. Note: Do not transfer cells with serum. If necessary, centrifuge a second time. Plasma (soluble C5b-9, C3c, Ba, Bb, FD, Properdin, C5, FB, FH, FI levels) Collection Protocol (minimum volume: 2ml) *PLEX treatments will affect plasma tests, please wait ~14 days after PLEX to draw samples: 1. Follow standard phlebotomy techniques to collect at least 6 cc of whole blood drawn in a lavender-top (EDTA) vacutainer tube. 2. Centrifuge at room temperature immediate after blood draw (1000 x g for 10 minutes). 3. Label "Plasma" or "Lavender-top" on clean screw top-tube(s). 4. Pipette cell-free supernatant (at least 2 mL) to each labeled tube (s). 5. Place the tube immediately at -80°C (or on dry ice). Sample must remain deep frozen. Note: Do not transfer cells with plasma. If necessary, centrifuge a second time. Plasma (ADAMTS-13 Activity/Inhibitor) Collection Protocol (minimum volume: 0.5ml): 1. Follow standard phlebotomy procedure to collect blood in 1. Following centrifugation, transfer the top two-thirds of the buffered sodium citrate (light blue-top, 3.2%) plastic tubes plasma layer into a new plastic tube. (available in 4.5 mL, 2.7 mL or 1.8 mL full draw tubes). Re-centrifuge the collected plasma at 1500 to 1800 x g 2. 2. After collection, invert the tube gently 5 to 6 times. with the brake off for an additional 15-20 minutes to 3. Label "Citrate Plasma" or "Blue-top" on clean cryovial screwremove any red cells or platelets. top tubes. 3. Transfer the top two-thirds of the plasma into the 4. Store the blue-top tube upright at room temperature until previously labeled cryovials, taking care not to disturb any centrifugation. Samples should be centrifuged between 15 cells at the bottom of the tube. to 60 minutes after blood collection for best results. 7. Place the tube immediately at -80°C (or on dry ice). 5. Re-mix the blood sample immediately prior to centrifugation Sample must remain deep frozen. by gently inverting the tube 5 to 6 times. 6. Centrifuge blood sample at room temperature in a horizontal Note: if the sample arrives at room temperature a new rotor (swinging bucket rotor) for 15-20 minutes at 1500 to sample will be required. 1800 x g with the brake off. Ship all samples to: Serum & Plasma Shipping Requirements: Serum and plasma must be frozen and shipped OVERNIGHT with a Dr. Richard Smith minimum of 3 kg (or 6 lbs) of dry ice. Molecular Otolaryngology & Renal Research Laboratories Cryovials should be put in zip lock bags and completely covered in dry ice to keep the sample frozen until it arrives in the lab. The University of Iowa 285 Newton Rd., 5270 CBRB Delivery: Monday-Friday. NO WEEKEND DELIVERIES lowa City, IA 52242-1078 Thawed OR unlabeled samples will be REJECTED for testing. Phone: 319-335-6623



Page 4 of 5

Ordanyngology & Renal Research Luboratories / The AETT COARCE Email: <u>morl@uiowa.edu</u> • <u>https://morl.lab.uiowa.edu</u> CLIA: 16D0966193					
Complement Panel tests offered by the MORL:	Test Code:				
C3 Glomerulopathy Complement Panel (serologies for DDD, C3GN), Serum and Plasma - CH50, APFA, C3b Deposition Assay, FHAA, FBAA, Fluid Phase Activity Assay-IFE, Nephritic Factors (C3Nef-C3CSA, C5Nef-C3CSAP, C4Nef), C3, C3c, C4, FB, Ba, Bb, FD, Properdin, C5, Soluble C5b-9, FH and FI levels					
aHUS (complement-mediated TMA Functional Panel (serologies for TTP, aHUS, HUS), Serum and Plasma - CH50, APFA, C3b Deposition, FHAA, FBAA, Fluid Phase Activity-IFE, C3, C3c, C4, FB, Ba, Bb, FD, Properdin, C5, Soluble C5b-9, FH and FI levels	aHUS-FP				
Autoantibody Panel, Serum - FHAA, FBAA, Fluid Phase Activity-IFE, Nephritic Factors (C3Nef-C3CSA, C5Nef- C3CSAP, C4Nef)	AAP				
Complement Biomarker Panel, Plasma - C3, C3c, C4, FB, Ba, Bb, FD, C5, Properdin levels, soluble C5b-9, FH and FI levels	СВР				
Complement Pathway Activity Panel, Serum - CH50, APFA, C3b Deposition Assay	СРАР				
Autoantibodies to Complement Components	Test Code:				
Fluid Phase Activity Assay, Serum (IFE)	07FPA				
FH Autoantibody, Serum (ELISA)	07FHAA				
FB autoantibody, Serum (ELISA)	07FBAA				
C3Nef, Serum (Hemolytic)	06C3NEF				
C5Nef, Serum (Hemolytic)	06C5NEF				
C4Nef, Serum (Hemolytic)	06C4NEF				
Functional Assays of Complement Activity - Pathways	Test Code:				
CH50, Serum (Liposome-based method)	07CH50				
Alternative Pathway Functional Assay (APFA), Serum (ELISA)	06APFA				
C3b Deposition Assay (Hemolytic)	01C3BDA				
Complement Protein Biomarkers (including split products)	Test Code:				
C3 Level, Serum (Turbidmetry)	07C3L				
C3c Level, Plasma (ELISA)	06C3CL				
C4 Level, Serum (Turbidmetry)	07C4L				
FB Level, Plasma (ELISA)	07FBL				
Ba Level, Plasma (ELISA)	06BAL				
Bb Level, Plasma (ELISA)	06BBL				
FD Level, Plasma (ELISA)	06C5L				
Properdin Level, Plasma (ELISA)	06PL				
C5 Level, Plasma (ELISA)	06C5L				
Soluble C5b-9, Plasma (ELISA)	06SMAC				
FI Level, Plasma (ELISA)	07FIL				
FH Level, Plasma (ELISA)	06FHL				
ADAMTS-13	Test Code:				
ADAMTS-13 Activity (a la carte only), Citrate Plasma (FRET)					
ADAMTS-13 Activity with reflex to Inhibitor Assay (if activity is <25%), Citrate Plasma (FRET)	01ATS13RFX				
Genetic Tests Offered by the MORL:	Test Code:				
Genetic Renal Panel: NGS + MLPA (CNVs) for Complement-Mediated Kidney Disease	GRP08				
MLPA (CFH-CFHR5): Multiplex Ligation Dependent Probe Amplification	MLPA02				

MORL – Kidney Testing Requisition Form - Please see page 2 for Tests Offered

Specimen and shipping requirements along with CPT codes and prices can be found on our website: https://morl.lab.uiowa.edu.



DISCLAIMER:

This request to order molecular diagnostic tests from the MORL certifies to the MORL that the ordering healthcare provider has obtained informed consent from the patient as required by applicable state or federal laws for each test ordered, that the ordering healthcare provider has authorization from the patient permitting the MORL to report results for each test ordered to the ordering physician, and that the ordering healthcare provider assumes responsibility for providing the patient with all associated guidance and counseling regarding the test results.

ALL requested information must be provided, or testing will not be performed

Patient information:	Patient date of birth and gender			
	Patient ethnicity and race			
	Patient's clinical information and family history of kidney disease			

We request extensive patient demographic and clinical information. This information is <u>required</u> as it is very valuable in the interpretation of your patient's results.

Specimen information:	Patient identifiers (full name, date of birth, sex, and medical record number) Date of collection <u>Sample type – frozen samples must be CLEARLY LABELED as either serum or plasma (and type, EDTA or Citrate)</u> Ordering physician
Billing information:	We will <u>NOT</u> bill insurance, Medicare, or patient directly. Institutional billing accepted. Visa and Master Card accepted. Personal checks <u>NOT</u> accepted. Please include contact information including phone & fax number for billing questions.
Reporting Information:	Because of confidentiality issues, reports will only be released to the individual indicated on the page 1 of the testing requisition form.
Research Participation:	If your patient's genetic and functional testing results are inconclusive, they may qualify for research studies on complement-mediated renal diseases that are ongoing at the MORL. If you would like your patient to be considered for this opportunity, please contact Amy Weaver at <u>amy-weaver@uiowa.edu</u> .

IMPORTANT INFORMATION FOR PHYSICIAN OR GENETIC COUNSELOR:

DNA tests may detect an abnormality. Detection methods are greater than 99% accurate. Many of these tests are relatively new. The analysis and interpretation represent our best knowledge and understanding of the genetics of these diseases.

There is a small possibility that a test may not work properly, or an error may occur. You may be asked for an additional sample if it is felt that confirmatory testing is needed.

An error in diagnosis may occur if incorrect information is provided with the sample.

Kidney diseases are complex disorders and penetrance of a phenotype (the degree of kidney disease, for example) may be variable. Research to determine whether a genotype-phenotype correlation exists is ongoing.

Because of the complexity of DNA testing, results should be discussed with a genetic counselor or physician.

Note: Kidney diseases are very complex disorders. This complexity means that variants in many different genes can lead to kidney disease. It is possible that no variants will be detected in the variant screens (the genes) you have requested.